



Docket No.: 5051-451ip

PATENT

**IN THE UNITED STATES PATENT AND TRADEMARK OFFICE**

In Re: Yuehua Li, et al.

Confirmation No.: 8515

Serial No.: 09/914,020

Group Art Unit: 1635

Filed: December 31, 2001

Examiner: Janet L. Epps-Ford

For: METHODS AND COMPOSITIONS FOR ALTERING MUCOUS SECRETION

## **Appendix A**

**IN THE UNITED STATES PATENT AND TRADEMARK OFFICE**

Atty. Docket No: 5051-4511P

In re patent application of

Yuehua LI *et al.*

Group Art Unit: 1635

Serial No. 09/914,020

Examiner: Janet L. Epps-Ford

Filed: December 31, 2001

For: METHODS AND COMPOSITIONS FOR ALTERING MUCUS SECRETION

**DECLARATION UNDER 37 C.F.R. §1.132**

I, Indu Parikh, declare that:

1. I hold the position of President and Chief Scientific Officer with BioMarck Pharmaceuticals. I am a biochemist and have worked in a variety of areas including protein/peptide chemistry from 1966 to the present. A copy of my Curriculum Vitae is appended hereto as Exhibit 1.

2. I have read and understood the rejections based on the alleged lack of written description and alleged lack of enablement for MARCKS derived peptides other than SEQ ID NO:1 (MANS peptide, Myristoylated N-terminal Sequence) in the above-captioned application, mailed on June 6, 2005

3. In response to these rejections, I provide the results of experiments that were performed at my request showing that other myristoylated (MA) peptides of the N-terminal sequence of the MARCKS protein inhibit mucus hypersecretion in mammalian airways. Experiments were designed to test whether these peptides inhibit hypersecretion of mucin in murine airways *in vivo*. The well-established ovalbumin-sensitized mouse model of asthma was used in these studies as described in Eum *et al.*, *Proc. Natl. Acad. Sci. USA* 92: 12290-12294, 1995 (a copy is attached as Exhibit 2). As a negative control, the RNS (Randomized N-terminal Sequence, also called BIO-124R) peptide was tested along with the other peptides. Specifically, BP2 mice (or Balb/C mice), aged 6-8 wks, were immunized subcutaneously twice at weekly intervals with 1µg of ovalbumin. After 14 days of sensitization, the animals were exposed to aerosolized ovalbumin, which causes a pronounced goblet cell hyperplasia. At the 72 hr time point, the secretagogue, methacholine (60 mM) was delivered using a Boxco system nebulizer providing a fine

aerosol for 90 seconds. Fifteen minutes prior to the secretagogue challenge, 50  $\mu$ L of the test peptide, at 3 different concentrations (10  $\mu$ M, 100  $\mu$ M, or 140  $\mu$ M), was administered to the mice by intratracheal route. The RNS peptide was tested at the highest dose level only (50  $\mu$ L of 140  $\mu$ M solution). The various controls used in these experiments included (a) untreated saline group, (b) OVA sensitized group in absence of any peptide, and (c) group in which RNS peptide was used as a control. Each experiment was carried out in 6 mice per group, and each set of experiments was repeated 3 times. To test strain-to-strain variations, the above experiment was repeated in Balb/C mice under similar protocol. Following the methacholine challenge, the animals were sacrificed and bronchoalveolar lavage (BAL) performed on 5 animals per group for analysis of secreted mucin by ELISA method and the resulting data normalized to mg protein.

4. The following controls and peptides of the N-terminal sequence of the MARCKS protein were tested in these experiments. The data is presented in Exhibit 3 and show the results of those myristoylated peptides tested at 100  $\mu$ M concentrations.

- a. Saline = untreated saline group (control);
- b. Ova = OVA sensitized group in absence of any peptide (control);
- c. BIO-124R = RNS peptide (control);
- d. BIO-124 = MANS peptide, Myristoylated 1-24 amino acid sequence of MARCKS protein.
- e. BIO-120 = Myristoylated amino acid sequence 1-20 of MARCKS protein;
- f. BIO-116 = Myristoylated amino acid sequence 1-16 of the MARCKS protein;
- g. BIO-112 = Myristoylated amino acid sequence 1-12 of MARCKS protein; and
- h. BIO-110 = Myristoylated amino acid sequence 1-10 of the MARCKS protein.

5. The results, as presented in Exhibit 3, demonstrate that the above peptides, designated BIO-124, BIO-120, BIO-116, BIO-112, and BIO-110, show between 66 to 77 % inhibition of mucin hypersecretion in the mouse model of asthma as compared to the Ova control. These data support the enablement and written description of the present invention for fragments from the N-terminal sequence of the MARCKS protein (SEQ ID NO: 3) including MANS peptide.

6. I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further, that these statements are made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issuing thereon.

September 6, 2005  
Date

  
Indu Parikh, Ph.D.

253576 v1/RE

Attorney Docket No.: 5051-451ip

PATENT

**IN THE UNITED STATES PATENT AND TRADEMARK OFFICE**

In Re: Yuehua Li, et al.

Confirmation No.: 8515

Serial No.: 09/914,020

Group Art Unit: 1635

Filed: December 31, 2001

Examiner: Janet L. Epps-Ford

For: METHODS AND COMPOSITIONS FOR ALTERING MUCOUS SECRETION

**Exhibit 1**

## EXHIBIT 1

INDU PARIKH, Ph.D.

4364 South Alston Ave

Durham, NC 27713

Tel: 919-768-1335

Fax: 919-361-2290

E-Mail: [iparikh@cato.com](mailto:iparikh@cato.com)

---

### PROFESSIONAL SUMMARY:

Over 30 years of industrial drug discovery and drug development experience in design and discovery of new products and concepts in the pharmaceutical areas including controlled drug delivery systems, endocrinology, osteoporosis, mucosal diseases of the gastrointestinal tract, mucus hypersecretory pulmonary diseases, and proteins/peptides as therapeutic agents. Productive innovator and R&D leader with record of turning good ideas into profitable business opportunities and advancement of inventions to clinical development and to marketable stage.

#### Strengths:

- Strongly diversified technical background with business acumen
- Practical, hands-on, goal oriented approach to problem solving
- Experience in business development
- Organizational, communication and project management skills

### PROFESSIONAL EXPERIENCE:

2002-	<b>Co-Founder, President &amp; CSO</b> BioMarck Pharmaceuticals, Ltd., Durham, North Carolina
1997-2000	<b>Co-Founder, Chief Scientific Officer and Vice President</b> RTP Pharma Inc. Ile des Soeurs, Verdun, QC, Canada
1990-1997	<b>Co-Founder and Vice President for Research &amp; Development</b> Research Triangle Pharmaceuticals Ltd. Durham, NC 27713
1986-1990	<b>Director</b> Division of Biochemistry and Biotechnology Glaxo Inc. Research Triangle Park, NC
1986-1988	<b>Director of Extramural Research</b> Glaxo Inc. Research Triangle Park, NC
1982-1986	<b>Head</b> Molecular Biology Department Burroughs Wellcome Co. Research Triangle Park, NC
1977-82	<b>Associate Head</b> Molecular Biology Department Burroughs Wellcome Co. Research Triangle Park, NC
1975-77	<b>Group Leader</b> Molecular Biology Department Burroughs Wellcome Co., RTP, NC.

## **EXHIBIT 1**

1970-75 **Assistant Professor of Pharmacology & Experimental Therapeutics and Assistant Professor of Medicine**  
Johns Hopkins University. School of Medicine, Baltimore, MD

1968-70 **Visiting Scientist**  
NIAMD, NIH, Bethesda, Maryland (Dr. Christian B. Anfinsen)

1966-68 **Postdoctoral Fellow**  
Weizmann Institute of Science, Rehovot, Israel (Prof. Ephraim Katzir)

1965-66 **Research and Teaching Associate**  
University of Zurich, Zurich, Switzerland

**EDUCATION:** Ph.D. University of Zurich, Zurich, Switzerland (1965)

### **PROFESSIONAL MEMBERSHIPS:**

American Society of Biochemistry and Molecular Biology  
American Society of Pharmacology and Experimental Therapeutics  
Control Release Society  
American Assoc. of Pharmaceutical Scientists  
International Society for Biorecognition Technology  
National Institute of Health Alumni Association

### **HONORS:**

Monsanto Postgraduate Fellowship, 1961-63  
Geigy Postgraduate Fellowship, 1963-65  
Weizmann Institute Visiting Fellowship, 1966-68  
Executive Vice President and Member, Board of Directors, Alopecia Areata Research Foundation, 1983-1990; 1991-1996  
Member, Governing Council, Internat. Society for Biorecognition Technology, 1984-90  
Treasurer, International Society for Biorecognition Technology, 1984-89  
Honorary Life Member, Society of Biological Chemistry of India, 1989-  
Honorary Member, National Academy of Sciences, India, 1989-

### **INDUSTRIAL AFFILIATIONS:**

Member of Scientific/Corporate Advisory Boards of:  
Bioxy International, 1989-1999  
Protein Delivery Inc., (Nobex Pharmaceuticals) 1991-1997  
Biomedica, Inc., 1989-1997  
Cibus Pharmaceuticals, 1992-1996  
Medicago, Inc., 1999-present  
Bioinfinix, Inc., 1999-2002  
RTP Pharma Inc., 2000-2002  
PhytoMyco Research Corp., 2002-present

## EXHIBIT 1

### EDITORIAL BOARDS:

Journal of Solid-Phase Biochemistry, 1975-80  
Journal of Applied Biochemistry and Biotechnology, 1980-86  
Pharmacology and Drug Development, 1979-86

### ACADEMIC APPOINTMENT:

Adjunct Professor, Pediatric Endocrinology, Department of Medicine, University of North Carolina School of Medicine at Chapel Hill, NC, 1987-1992.

### MAJOR ACHIEVEMENTS:

#### **Research Triangle Pharmaceuticals Ltd. (RTP Pharma Inc.), 1990-2000**

The Company was founded on the yet to be exploited concepts in delivery of water insoluble drugs. The RTP's proprietary drug delivery technology for water insoluble drugs, DD™, was explored and employed for oral, injectable, aerosol and topical delivery of more than 100 different drugs. Two of these drugs are currently at NDA stage. The Company was initially founded in North Carolina in 1990 and later funded by a group of Quebec based institutional investors and was relocated to Montreal in 1996. RTP Pharma was acquired by SkyePharma in 2002.

The following is a few of the highlights of my contributions to RTP Pharma.

- Co-founded the Company.
- Established a data base for water insoluble drugs, performed feasibility studies, developed clientele for contract research, expanded intellectual property base for the company's proprietary drug delivery technology.
- Wrote business plan, actively participated in fund raising activities in U.S. and Canada.
- After relocation to Canada (December 1996), helped recruiting additional staff at all levels, including four VPs (Medical, Process Dev., Business Dev., and CFO), prioritized projects, and provided overall day to day R&D direction, successful in gaining R&D contracts from five large pharmaceutical companies, and actively participated in other business development activities as needed.
- Helped select drug candidates and implementation for in-house development. Lead two of these drugs to NDA stage.
- Performed proof-of-concept key experiments for a technology based on differentiation of islet precursor cells as a means to treat Type II diabetes. This technology was then spun off as a subsidiary company, Waratah Pharmaceuticals, Inc. Waratah went public in September 2000 with Canadian Small Cap CDNX exchange. This was CDNX's top selling IPO for the year 2000.

#### **Director, Biochemistry & Biotechnology Division, Glaxo Inc., 1986-1990**

The Division was active in goal oriented discovery research in biopharmaceuticals. Collaboration with University investigators was encouraged in selected areas. The Division consisted of five departments namely, Molecular Biology, Enzymology, Protein Chemistry,



## EXHIBIT 1

### Biochemical Pharmacology, and Hybridoma Technology.

- Molecular Biology Department: An important function of this Department was to express desired proteins by recombinant technology for detail biochemical and pharmacological studies. Expression systems like E. coli, mammalian cells, and yeast were routinely used. Some of the proteins expressed included:
  - Lipocortin I and II ■ tissue inhibitor of metalloproteases ■ EGF ■ hSOD
  - extracellular region of CD4 and ■ a fibrinolytic enzyme ■ development of vector systems to enhance the yield of cloned proteins ■ site specific mutagenesis, ■ gene regulation studies with lipocortin I and two lymphokines.
- Enzymology Department: Major efforts in the rational drug development based on designing specific inhibitors of selected metabolic enzymes. The major activities of the department included: two fibrinolytic enzymes, metalloproteases including collagenase, and 5 $\alpha$ -reductase, and aromatase.
- Protein Chemistry Section: The section primarily provided purification and scale-up services for proteins generated by recombinant labs.
- Biochemical Pharmacology: This section investigated primarily the mucosal diseases of GI tract including gastric ulcer, atrophic gastritis, IBS, etc.: helped develop highly sensitive quantitative digital video endoscopy technology. In addition, developed a rat model for osteoporosis in collaboration with a local university.
- Hybridoma Center: The primary charter of this section was to provide monoclonal and polyclonal antibodies for research and for cloning purposes for the company.
- Miscellaneous:
  - Helped establish Glaxo's U.S. R&D operation from ground zero. Helped R&D division grow from two employees and zero sq. ft to about 450 staff and more than 700,000 sq. ft. of lab space, helped establish and prioritize selected therapeutic areas for the U.S. operation. Helped in-licensing activities for the U.S. Corporation.

### Head, Molecular Biology Department, Burroughs Wellcome Co., 1975-1986

The research emphasis in the Molecular Biology Department was centered on hormonal action, hormone receptors, and signal transduction mechanisms. Among the hormones investigated included, EGF, insulin, estrogen, progesterone, IGF-I, interferon, and opiate. Beside investigating enzymes like adenylate cyclase, phospholipase C, Phospholipase A2 and G-proteins, the Department also provided a general screening services to the Company for various receptors and selected enzymes.

Both functionally and administratively I was also responsible for The Wellcome Diagnostics Research. In addition to the administrative and scientific responsibilities of the Department, the following can be listed as major achievements of my own research activities.

- Established, organized, and conducted Therapeutic Advisory Panel.
- Purified and characterized estrogen receptors.
- Generated monoclonal antibodies to estrogen receptor (1977).

## EXHIBIT 1

- Devised and patented commercially viable diagnostic assay for estrogen receptor.
- Established and patented a novel homogeneous enzyme immunoassay.
- Discovered and patented a non-absorbable antidiarrheal drug.
- Acted as Project Leader for preclinical work on the novel antidiarrheal drug.
- Cloned and expressed Cu-Zn Superoxide Dismutase (hSOD).
- Purified  $\alpha$  interferon (1976)
- Actively evaluated external licensing candidates for both Burroughs Wellcome Co. and for Wellcome Diagnostics Ltd.

## BIBLIOGRAPHY:

1. Mabry, T.J., Wyler, H., Sassu, G., Mercier, M., Parikh, I. and Dreiding, A.S. Die Struktur des Neobetamidins. Über die Konstitution des Randfarbstoffes Betanin. Helvetica Chimica Acta 45, 640-647, 1962.
2. Mabry, T.J., Wyler, H., Parikh, I. and Dreiding, A.S. The Conversion of Betanidin and Betanin to Neobetamidin. Tetrahedron 23, 3111-3127, 1967.
3. Badgett, B., Parikh, I. and Dreiding, A.S. Totalsynthese Eines Neobetamidin Derivates und des Neobetamins. Helvetica Chimica Acta 53, 433-448, 1970.
4. Parikh, I. and Omenn, G.S. Modification of Staphylococcal Nuclease with Nitrophenylsulfenyl Halides. Biochemistry 10, 1173-1177, 1971.
5. Parikh, I., Corley, L. and Anfinsen, C.B. Semisynthetic Analogues of an Enzymatically Active Complex Formed Between Two Overlapping Fragments of Staphylococcal Nuclease. J. Biol. Chem. 246, 7392-7397, 1971.
6. Cuatrecasas, P. and Parikh, I. Adsorbents for Affinity Chromatography. Use of N-Hydroxysuccinimide Esters of Agarose. Biochemistry 11, 2291-2299, 1972.
7. Parikh, I. and Cuatrecasas, P. Affinity Chromatography. Vox Sanguinis 23, 141-146, 1972.
8. Krug, F., Parikh, I., Illiano, G. and Cuatrecasas, P.  $\beta,\gamma$ -Methylene-Adenosine-5'-Triphosphate - A Competitive Inhibitor of Adenylate Cyclase in Fat and Liver Cell Membranes. J. Biol. Chem. 248, 1203-1206, 1973.
9. Krug, F., Parikh, I., Illiano, G. and Cuatrecasas, P.  $\beta,\gamma$ -Methylene-Adenosine-5'-Triphosphate - Effects of a Competitive Inhibitor of Adenylate Cyclase on Cyclic AMP Accumulation and Lipolysis in Isolated Fat Cells. Biochem. Biophys. Res. Commun. 50,

## EXHIBIT 1

985-991, 1973.

10. Parikh, I. and Cuatrecasas, P. Enzyme Immobilization and Adsorbents with Biologic Specificity. In: Enzyme Therapy in Genetic Diseases, D. Bergsma (Ed.), Williams and Wilkins Co., Baltimore, MD., Vol. IX 43-45, 1973.
11. Sica, V., Nola, E., Parikh, I., Puca, G.A. and Cuatrecasas, P. Purification of Oestradiol Receptors by Affinity Chromatography. Nature New Biology 244, 36-39, 1973.
12. Sica, V., Parikh, I., Nola, E., Puca, G.A. and Cuatrecasas, P. Affinity Chromatography and Purification of Estrogen Receptors. J. Biol. Chem. 248, 6543-6558, 1973.
13. Cuatrecasas, P., Parikh, I. and Hollenberg, M.D. Affinity Chromatography and Structural Analysis of Vibrio cholerae Enterotoxin-Ganglioside Agarose and the Biological Effects of Ganglioside-Containing Soluble Polymers. Biochemistry 12, 4253-4264, 1973.
14. Parikh, I. and Cuatrecasas, P. Substrate Analog Competitive Inhibitors of Human Renin. Bioch. Biophys. Res. Comm. 54, 1356-1361, 1973.
15. Parikh, I. and Cuatrecasas, P. Affinity Chromatography. In: Reactions on Polymers, J.A. Moore (ed), D. Reidel Publishing Co., Boston, Mass., 340-343, 1973.
16. Cuatrecasas, P., Tell, G.P.E., Sica, V., Parikh, I. and Chang, K.-J. Noradrenaline Binding and the Search for Catecholamine Receptors. Nature 247, 92-97, 1974.
17. March, S.C., Parikh, I. and Cuatrecasas, P. A Simplified Method for Cyanogen Bromide Activation of Agarose for Affinity Chromatography. Anal. Biochem. 60, 149, 152, 1974.
18. March, S.C., Parikh, I. and Cuatrecasas, P. Affinity Chromatography - Old Problems and New Approaches. In: Immobilized Biochemicals and Affinity Chromatography, R.B. Dunlap (Ed.), Plenum Press, New York 3-14, 1974.
19. Grebner, E.E. and Parikh, I. An Affinity Adsorbent for N-Acetyl- $\alpha$ -D-Hexosaminidase. Biochem. Biophys. Acta 350, 437-441, 1974.
20. Parikh, I., March, S. and Cuatrecasas, P. Topics in the Methodology of Substitution Reactions with Agarose. Methods of Enzymology 34 77-102, 1974.
21. Parikh, I. and Cuatrecasas, P. Ganglioside-Agarose and Cholera Toxin. Methods in Enzymology 34, 610-619, 1974.
22. Parikh, I., Sica V., Nola, E., Puca, G.A. and Cuatrecasas, P. Estrogen Receptors. Methods

## EXHIBIT 1

in Enzymology 34, 670-688, 1974.

23. Cuatrecasas, P. and Parikh, I. Insulin Receptors. Methods in Enzymology 34, 653-670, 1974.
24. Quaroni, A., DiCapua, E., Parikh, I. and Semenza, G. Affinity Labeling of Sucrose-Isomaltase Complex from Rabbit Small Intestine. Experientia 30, 690, 1974.
25. Parikh, I. and Cuatrecasas, P. Principles and Applications of Affinity Chromatography. In: Methods in Protein Separation, N. Catsimpoolas (Ed.), Plenum Press, New York, pp. 255-276, 1975.
26. March, S. C., Pasternak, G.W., Parikh, I., Snyder, S.H. and Cuatrecasas, P. Macromolecular Naloxone: A Long-Acting Opiate Antagonist. In: Problems of Drug Dependence, 607-11, 1975.
27. March, S., Parikh, I. and Cuatrecasas, P. Purification of Hormones and Other Plasma Components by Affinity Chromatography. In: Hormones in Human Plasma, Vol. I, H.N. Antoniades (Ed.), Harvard University Press, pp. 32-50, 1976.
28. Pasternak, G., March, S., Parikh, I., Snyder, S.H. and Cuatrecasas, P. Macromolecular Naloxone: A Novel Class of Long-Acting Polymer Drug. Life Sciences 18, 977-982, 1976.
29. Parikh, I., MacGlashan, D.C. and Fenselau, C. Immobilized Glucuronyl Transferase for the Synthesis of Conjugates. J. Med. Chem. 19, 296-299, 1976.
30. Sica, V., Nola, E., Puca, G.A., Cuatrecasas, P. and Parikh, I. Purification of Estrogen Receptor by Affinity Chromatography. In: Receptors and Mechanism of Action of Steroid Hormones, J.R. Pasqualini, (Ed.), pp. 85-108 Marcel Dekker, Inc., New York, 1976.
31. Fenselau, C., Pallante, S. and Parikh, I. Solid-phase Synthesis of Drug Glucuronides by Immobilized Glucuronyltransferase. J. Med. Chem. 19 679-683, 1976.
32. Sica, V., Cuatrecasas, P., Nola, E., Parikh, I. and Puca, G.A. Purification of Estrogen Receptor by Affinity Chromatography. In: Mod. Pharmacol. Toxicol. 8, 85-108, 1976.
33. Parikh, I. and Cuatrecasas, P. Affinity Chromatography in Immunology. In: Immunochemistry of Proteins, M.Z. Attasi (Ed.) Vol. 2, pp. 1-44. Plenum Press, New York, 1977.
34. Parikh, I. and Slovin, S.F. An Introduction to Affinity Chromatography. In: Cancer Biology IV: Differentiation and Carcinogenesis, C. Borek, C.M. Fenoglio and D.W. King

## EXHIBIT 1

- (Eds.), Stratton Intercontinental, 1977.
35. Anderson, W.L., Shechter, Y. and Parikh, I. Quantitation of Methionyl Peptides in Nanomole Quantities by a Fluorometric Method. Anal. Biochem. 91, 481-489, 1978.
  36. Parikh, I. Isolation of Hormone Receptors by Affinity Chromatography. In: Affinity Chromatography, O. Hoffmann-Ostenhof (Ed.), Pergamon Press, 1978 pp. 319-327.
  37. Parikh, I., Anderson, W.L. and Neame, P.J. Identification of High Affinity Estrogen Binding Sites in Calf Uterine Membranes. J. Biol. Chem. 255, 10266-10270, 1980.
  38. Parikh, I. and Cuatrecasas, P. Hormone-Receptor Interactions in Homogeneous and Heterogeneous Systems: Relationship to Affinity Chromatography. In: Affinity Chromatography and Related Techniques. T.C.J. Gribnau, et al(Eds.) Elsevier, 1981, pp.79-82.
  39. Nola, E., Molinari, A.M., Medici, N., Moncharmont, B., Piccoli, R., Puca, G.A., Parikh, I. and Cuatrecasas, P. Purification of the Native Form of the Oestradiol Receptor. In: Research on Steroids, Vol. 8, (Eds. A. Kloppe et al.) Academic Press, pp. 167-174, 1979.
  40. Moncharmont, B., Parikh, I., Puca, G.A. and Cuatrecasas, P. Effect of Sodium Molybdate on Cytosolic Estrogen Receptor. J. Steroid Biochem. 16, 361-368, 1982.
  41. Neame, P.J. and Parikh, I. Sepharose Immobilized Triazine Dyes as Adsorbents for Human Lymphoblastoid Interferon Purification. J. Appl. Biochem. Biotech. 7, 295-305, 1982.
  42. Moncharmont, B., Su, J.-L. and Parikh, I. Monoclonal Antibodies Against Estrogen Receptors: Interaction with Different Molecular Forms and Functions of the Receptor. Biochemistry 21, 6916-6921, 1982.
  43. Moncharmont, B. and Parikh, I. Binding of Monoclonal Antibodies to the Nuclear Estrogen Receptor in Intact Nuclei. Biochem. Biophys. Res. Comm. 114, 107-112, 1983.
  44. Chaiken, I.M., Wilcheck, M. and Parikh, I. (Eds.). Affinity Chromatography and Biological Recognition. Academic Press, New York, pp. 515, 1983.
  45. Moncharmont, B., Anderson, W.L., Rosenberg, B. and Parikh, I. Interactions of Estrogen Receptor of Calf Uterus with a Monoclonal Antibody: Probing of Various Molecular Forms. Biochemistry 23, 3907-3912, 1984.
  46. Moncharmont, B. and Parikh, I. Monoclonal Antibodies to Steroid Receptors. In: Receptors and Recognition (Ed. M.F. Greaves), Chapman & Hall, pp. 67-86, 1984.

## EXHIBIT 1

47. Parikh, I. and Cuatrecasas, P. Horizons in Affinity Chromatography and Its Application. Chem. & Eng. News. 63, 17-32, 1985.
48. Parikh, I. and Cuatrecasas, P. Affinity Chromatography. In: CRC Handbook of Clinical Chemistry, (M. Werner Ed.), CRC Press, 1981. Boca Raton, Florida, pp. 307-314, 1985.
49. Moncharmont, B., Parikh, I., Anderson, W.L., Su, J.-L., Massol, N., She, W.-M. and Lebeau, M.C. Applications of Monoclonal Antibodies to the Study of Estrogen Receptor Structure. In: Monoclonal Antibodies: Biological and Clinical Applications. (Eds. A. Pinchera, et al), Editrice Durtis Press, pp. 259-276, 1985.
50. Sar, M. and Parikh, I. Immunohistochemical Localization of Estrogen Receptor in Rat Brain and Pituitary with Monoclonal Antibodies. J. Steroid Biochem. 24, 497-503, 1986.
51. Parikh, I., Hilpert, H., Hermann, K. and Dreiding, A. Betenamin and Betalain Modelsubstanzen. Helv. Chim. Acta 69, 1588-1596, 1986.
52. Rajendran, K.G., Lopez, T. and Parikh, I. Estrogenic Effect of Phenol Red in MCF-7 Cells is Achieved Through Activation of Estrogen Receptor by Interacting with a Site Distinct from the Steroid Binding Site. Biochem. Biophys. Res. Comm. 142, 724-731, 1987.
53. Kull, F.C., Brent, D.A., Parikh, I. and Cuatrecasas, P. Chemical Identification of a Tumor-Derived Angiogenic Factor. Science 236, 843-845, 1987.
54. Marchetti, E., Querzoli, P., Moncharmont, B., Parikh, I., Bagni, A., Marzola, A., Fabris, G., and Nenci, I. Immunocytochemical Demonstration of Estrogen Receptors by Monoclonal Antibodies in Human Breast Cancer: Correlation with Estrogen Receptor Assays by Dextran Coated Charcoal Method. Cancer Research 47, 2508-2513, 1987.
55. Rajendran, K.G. and Parikh, I. Cellular Localization of Steroid Hormone Receptors by Immunocytochemistry. In: Receptor Mediated Hormone Antagonism, Walter de Gruyter, Publishers, Paris, 1987, pp. 469-496.
56. Lalwani, N.D., Alvares, K., Reddy, M.K., Reddy, M.N., Parikh, I., and Reddy, J.K. Peroxisome Proliferator-Binding Protein: Identification and Partial Characterization of Nafenopin- Clofibrac Acid-, and Clofibrate-Binding Proteins from Rat Liver. Proc. Natl. Acad. Sci. USA 84, 5242-5246, 1987.
57. Parikh, I., Rajendran, K.G., Su, J.-L., Lopez, T., and Sar, M. Are Estrogen Receptors Cytoplasmic or Nuclear? Some Immunocytochemical and Biochemical Studies. J. Steroid Biochem. 27, 185-192, 1988.

## EXHIBIT 1

58. Rajendran, K.G. and Parikh, I. Intracellular Localization of Estrogen Receptor. In: New Methods in Drug Research Vol 3 (Ed. A. Makriyannis), J. R. Prous Science Press, 1990, pp. 239-254.
59. Termanini, B., Nardi, R.V., Finan, T., Parikh, I. and Korman, L. Insulin-Like Growth Factor I Receptors in Rabbit Gastrointestinal Tract: Characterization and Autoradiographic Localization. Gastroenterology, 99, 51-60, 1990.
60. Anderson, J.J.B., Garner, S.C., Mar, M.H., Boass, N., Toverud, S.U. and Parikh, I. The Ovariectomized Lactating Rat as an Experimental Model for Osteopenia: Calcium Metabolism and Bone Changes. Bone and Mineral 11, 43-53, 1990.
61. Garner, S.C., Anderson, J.J.B., Mar, M.H. and Parikh, I. Estrogens Reduce Bone Loss in the Ovariectomized Lactating Rat Model. Bone and Minerals 15, 19-31, 1991.
62. Amarant, T., Burkhardt, W., LeVine, H., Arocha-Pinango, C.L. and Parikh, I. Isolation and Complete Amino Acid Sequence of two Fibrinolytic Proteases from the toxic Saturniid Caterpillar Lanomia achelous. Biochem. Biophys. Acta, 1079, 214-221, 1991.
63. Nardi, R.V., Guglietta, A. and Parikh, I. Epidermal Growth Factor. In: Handbook of Experimental Pharmacology: Pharmacology of Peptic Ulcer Disease, Vol. 99, pp. 37-54, Springer Verlag, 1991.
64. Burkhardt, W., Smith, G.F.H., Su, J.L., Parikh, I. and LeVine, H. Amino Acid Sequence Determination of Ancrod, the Thrombin-like  $\alpha$ -Fibrinogenase from the Venom of Akistrodon rhodostoma. FEBS, 297, 297-301, 1992.
65. Parikh, I. and Cuatrecasas, P. Affinity Chromatography: An Overview. In: Molecular Interactions in Bioseparations. (Ed. T. T. Ngo) pp. 3-13, Plenum Press, 1993.
66. Proia, A.D., Hirakata, A., McInnes, J.S., Scroggs, M.W. and Parikh, I. The effect of angiostatic steroids and  $\beta$ -cyclodextrin tetradecasulfate on corneal neovascularization in rat. Exp. Eye Res 57, 693-698, 1993.
67. Pace, S.N., Pace, G.W., Parikh, I., and Mishra, A.K. Novel Injectable Formulations of Insoluble Drugs. Pharmaceutical Technology 23, 116-134, 1999.
68. White, H.S., Woodhead, J.H., Wolf, H.H., Parikh, I., Clement, M., and Sofia, R.D. MicroParticle Felbamate Blocks Maximum Electroshock Following Intravenous Administration. Res. Comm. Biol. Psychology and Psychiatry 23, 1-7, 1998
69. Amarant, T., Nardi, R., and Parikh, I. Characterization of the Biological Activity of Human

## EXHIBIT 1

Epidermal Growth Factor. In preparation.

### ABSTRACTS:

1. Parikh, I. and Cuatrecasas, P. Purification of Cholera Toxin on Ganglioside-Agarose Affinity Adsorbents. Federation Proceedings 32, 559, 1973.
2. Sica, V., Nola, E., Puca, G.A., Parikh, I. and Cuatrecasas, P. Affinity Chromatography of Estrodiol Receptors of the Uterus. Federation Proceedings 32, 454, 1973.
3. Parikh, I. Substrate Analogous Inhibitors of Human Renin. Ninth International Congress of Biochemistry, Stockholm, Sweden, July 1973, p.98.
4. Parikh, I. and Cuatrecasas, P. Affinity Chromatography. 166th National Meeting of the American Chemical Society, Chicago, Illinois, August 26-31, 1973.
5. March, S.C., Parikh, I. and Cuatrecasas, P. Affinity Chromatography - Old Problems and New Approaches. Southeastern Regional Meeting of the American Chemical Society, Durham, North Carolina, August 1973.
6. Parikh, I., Agre, P. and Cuatrecasas, P. Isolation and Purification of E. coli Enterotoxin. Sixth International Congress of Pharmacology, Helsinki, Finland, July 20-25, 1975.
7. MacGlashan, D., Fenselau, C. and Parikh, I. Immobilization of UDP-Glucuronyl Transferase. Federal Proceedings 34, 812, 1975.
8. March, S., Pasternak, G., Parikh, I., Snyder, S. and Cuatrecasas, P. Pharmacodynamics of a Long Acting Opiate Antagonist in Mice: Macronaloxone. Clin. Rec. 23, 22A, 1975.
9. Keusch, G.T., Parikh, I. and Jacewicz, M. Affinity Chromatography: Purification of Shigella Toxin on a Sepharose-chitin Column. Am. Soc. Clin. Invest., January 1977.
10. Jacobs, S., Parikh, I. and Cuatrecasas, P. Purification of Insulin Receptors. Federation Proceedings 36(3), p. 710, 1977.
11. Lapetina, E.G., Chandrabose, K.A., Parikh, I. and Cuatrecasas, P. Platelet Aggregation Occurs in the Absence of Endoperoxides and Thromboxanes. Fed. Proc. 37(3), p. 259, 1978.
12. Hazum, E., Chang, K.-J., Parikh, I. and Cuatrecasas, P. Interaction of Iodinated Human (D-Ala<sup>2</sup>)-Endorphin and Opiate Receptors. Federation Proceedings 38(3), p. 457, 1979.
13. Shechter, Y., Jacobs, S., Chang, K.-J., Willingham, M., Pastan, I., Parikh, I., Cuatrecasas, P.



## EXHIBIT 1

- and Schlessinger, J. Mobility and Aggregation of Insulin and Epidermal Growth Factor (EGF) Receptors Complexes of the Plasma Membrane of Viable Fibroblasts. FASEB, Atlanta, Georgia, Federation Proceedings, June 1979.
14. Parikh, I. Steroid Hormone Receptor Systems. Edited by W.W. Leavitt and T.H. Clark, plenum Press, New York, 1979, 489 pp. (Book Review).
  15. Parikh, I., Anderson, W.L. and Neame, P. Evidence for a Membrane Associated Estradiol Receptor in Rat Uterus. Federation Proceedings 39(3), 980, 1980.
  16. Neame, P.J., Parikh, I. and Acton, R.T. The Potential of Triazine Dyes as Adsorbents for the Purification of Human Lymphoblastoid Interferon, 4th International Symposium on Affinity Chromatography and Related Techniques. Veldhoven, Netherlands, Abstract #B45, 1981.
  17. Parikh, I. Affinity Chromatography: Bioselective Absorption on Inert-Matrices by W.H. Scouten, John Wiley, New York, 1981, Applied Biochem. Biotech. (Book Review).
  18. Neame, P.J. and Parikh, I. Purification of Human Lymphoblastoid Interferon by Affinity Chromatography. Federation Proceedings 40(3), 4654, 1981.
  19. Parikh, I. and Cuatrecasas, P. Hormone-Receptors Interactions in Homogeneous and Heterogeneous Systems: Relationship to Affinity Chromatography. 4th Internat. Symposium on Affinity Chromatography and Related Techniques. Veldhoven, Netherlands, Abstract #L6, 1981.
  20. Moncharmont, B., Puca, G.A. and Parikh, I. Effect of Sodium Molybdate on Estrogen Receptor. Federation Proceedings 40, 1873, 1981.
  21. Moncharmont, B. and Parikh, I. Monoclonal Antibodies to Estrogen Receptors. UCLA Symposium. J. Cellular Biochem, Supplement 6, 163, 1982.
  22. Parikh, I. and Moncharmont, B. Monoclonal Antibodies to Estrogen Receptors. J. Steroid Biochem. 17, 101, 1982.
  23. Parikh, I. Targeted Drugs. Ed. E.P. Goldberg, John Wiley, New York, 1983. J. Med. Chem. 27, 101, 1984. (Book Review).
  24. Sar, M. and Parikh, I. Immuno-histochemical Localization of Estradiol Receptor in Rat Brain with Monoclonal Antibodies to Calf Cytosolic Receptor. 7th International Congress of Endocrinology, Quebec, Canada, p. 1468, 1984.
  25. Parikh, I. Monoclonal Antibodies: Studies on Various Molecular Forms of Estrogen

## EXHIBIT 1

Receptor. IV Pan American Biochemistry Congress, Buenos Aires, Argentina, Abstract #187, 1984.

26. Sar, M. and Parikh, I. Intercellular Localization of Estrogen Receptors in MCF-7 Cells with Monoclonal Antibodies. 67th Endocrine Soc. Meeting, Baltimore, Maryland, p. 289, 1985.
27. Parikh, I. Affinity Chromatography: A Practical Approach. Edited by P.D.G. Dean, W.S. Johnson and F.A. Middle, IRL Press, 1985. Nature 319, 629, 1986. (Book Review).
28. Parikh, I. Affinity Chromatography: Template Chromatography of Nucleic Acids. Edited by H. Schott, Dekker, 1985. Nature 319, 629, 1986. (Book Review).
29. Garner, S.C., Anderson, J.J.B. and Parikh, I. Estrogens Reduce Bone Loss in Ovariectomized Lactating Rats. J. Bone Min. Res., 4(1), 1989, Abstract #534.
30. Rajendran, K.G., Cortizo, A.M., Delacy, P. and Parikh, I. Evidence for Direct Effect of Steroid Hormones on Osteoblastic Cell Line UMR 106. Am. Soc. of Bone and Mineral Research, Montreal, Canada, 1989.
31. Garner, S.C., Anderson, J.J.B., Parikh, I. and Petrow, V. Nineteen Nor-Androstenediol Significantly Reduces Bone Loss in the Ovariectomized Lactating Rat. J. Bone Min. Res. 5(2), 1990, Abstract #406.
32. Anderson, J.J.B., Garner, S.C., Petrow, V. and Parikh, I. Skeletal Effects of Androstendiol in the Ovariectomized Lactating Rat. J. Bone Min. Res. 5(2), 1990, Abstract #405.
33. Haynes, D.H. and Parikh, I. Sustained Delivery of Indomethacin with a Novel Microcrystal Technology-Based Formulation. Am. Soc. Pharm. Exp. Ther. San Diego, CA p. 151, 1991.
34. Parikh, I., Watzek, W. and Haynes, D.H. Microcrystal Technology-Based sustained Delivery of Nifedipine in Rats. Am. Soc. Pharmacol. Exp. Ther. San Diego, p. 197, 1991.
35. McInnes, J., Hirakata, A., Scroggs, M.W., Parikh, I. and Proia, A.D. The Effect of Angiostatic Steroids and beta Cyclodextrins Tetradeca Sulfate on Corneal Neovascularization in the Rat. Invest. Ophthal. Visual Science, Annual Meeting, 1992, Sarasota, Florida, Abstract p. 1413.
36. Guglietta, A., Amarant, T., Parikh, I., Lesch, C. and Nardi, R.V. Effect of a Pharmacological Active Fragment of EGF on Gastric Acid Secretion in Rats and Monkeys. Gastroenterology, 4(2), A735, 1992.
37. Clement, M. A., Pugh, W. and Parikh, I. Tissue Distribution and Plasma Clearance of a Novel Microcrystal Fluribiprofen Formulation. Am. Soc. Pharmacology & Experimental

## EXHIBIT 1

- Therapeutics, National Meeting, Orlando, FL The Pharmacologist 34(3), 204, 1992.
38. McCulloch, W., Parikh, I., Colvin, O. M. and Paro, G. Development of I.V. Busulfan (Spartaject™-Busulfan). Proceedings of the Annual Meeting of European Group for Bone Marrow Transplantation, March 1994, Abstract # 185, p.116.
  39. Parikh, I. Delivery of Water Insoluble Drugs. Proceedings of the Annual Meeting of the Society for Industrial Microbiology, August, 1996, Abstract # S187, p.99.
  40. Parikh, I., Insoluble Drug Delivery Technology. Proceedings of the 1998 Conference on Pharmaceutical Science and Technology.
  41. Mercure, L., Chu, Q., and Parikh, I., Preclinical Assessment of a Novel Class of Immunoadjuvants. Proceedings of the First Annual Conference on Vaccine Research. 1998.
  42. Parikh, I. Affinity Chromatography: Lessons from the Past and Revelations for the Future. Proceedings of the 8<sup>th</sup> International Conference on Polymer Based Technology, Jerusalem, Israel, 1998.
  43. Carrier, A., Nguyen, X., Moussa, A., and Parikh, I. Rapid determination of paclitaxel in blood by HPLC with UV detection. Proceedings of the Annual AAPS Meeting, Abstract # 3530, page S-511, 1998.
  44. Bone, A.J., Lally, F.J., Lane, A., Parikh, I., and Barley, J. Stimulation of Islet Growth and Amelioration of Diabetes by Combination Therapy with TGF- $\alpha$  and Gastrin. Diabetologia, in preparation.

### PATENTS:

1. U.S. Patent No. 3,947,352 March 30, 1976  
Polysaccharide Matrices for Use as Adsorbents in Affinity Chromatography Techniques.
2. U.S. Patent No. 4,125,492 August 10, 1976  
Ganglioside Polysaccharide.

## EXHIBIT 1

3. British Patent No. 17749/78 May 4, 1978  
Method and Kit for Enzyme Immunoassay of Biological Substances.
4. U.S. Patent No. 4,225,487 April 13, 1978  
Biological Effects of Ganglioside Containing Polymers.
5. U.S. Patent No. 4,411,832 July 27, 1981  
Polysaccharide Matrices Comprising Macromolecular Spacers.
6. U.S. Patent No. 4,298,685 November 3, 1981  
Diagnostic Reagent Kit for Estrogen Receptor.
7. U.S. Patent No. 07/343,782 April 26, 1989  
Novel DNA Sequences Which Encode Ancrod-Like Polypeptides and Compositions  
Comprising the Expression Products.
8. U.S. Patent No. 5,434,135 July 18, 1995  
Growth Factor Composition, Preparation and Use.
9. U.S. Patent No. 5,785,975 July 28, 1998  
Novel Adjuvant Compositions and Vaccine Formulations.
10. U.S. Patent No. 5,660,858 August 26, 1997  
Cyclosporine Emulsions.
11. U.S. Patent No. 5,922,355 July 13, 1999  
MicroParticle Formulations of Water Insoluble Drugs
12. World Patent No. WO 99-49848 October 1999  
Anticancer Compositions.
13. World Patent No. WO 99-29300 June 1999  
Self-Emulsifying Fenofibrate Formulations.
14. U.S. Patent No. 6,086,376 July 2000  
MicroParticle Formulations for Aerosol Application.
15. World Patent No. WO 00-30616 June 2000  
Dispersible Phospholipid Stabilized Microparticles.
16. World Patent No. WO 00-30615 June 2000  
Method of Preparing Stable Suspension of Insoluble Microparticles.

## EXHIBIT 1

17. U.S. Patent No. 6,228,399 May 8, 2001  
Composition and Method of preparing MicroParticle  
of water Insoluble Substances.
18. U.S. Patent No. 6,465,016 October 15, 2002  
Cyclosporine Particles
19. U.S. Patent No. 6,558,952 May 6, 2003  
Treatment for diabetes
20. U.S. Patent No. 6,534,088 March 18, 2003  
Fibrate-Statin Combinations with Reduced Fed-Fasted Effects in  
Human Subjects.
21. U.S. Patent No. 6,696,084 February 24, 2004  
Spray Drying Process and Compositions of Fenofibrate.
22. World Patent No. WO 94/00015  
Biocidal Antimetabolite Compositions.

June 2003

\* \* \*

Attorney Docket No.: 5051-451ip

PATENT

**IN THE UNITED STATES PATENT AND TRADEMARK OFFICE**

In Re: Yuehua Li, et al.

Confirmation No.: 8515

Serial No.: 09/914,020

Group Art Unit: 1635

Filed: December 31, 2001

Examiner: Janet L. Epps-Ford

For: METHODS AND COMPOSITIONS FOR ALTERING MUCOUS SECRETION

**Exhibit 2**

# Eosinophil recruitment into the respiratory epithelium following antigenic challenge in hyper-IgE mice is accompanied by interleukin 5-dependent bronchial hyperresponsiveness

(asthma/airway eosinophilia/mouse model)

SEOK-YONG EUM\*, SOLOMON HAILÉ†, JEAN LEFORT\*, MICHEL HUERRE†, AND B. BORIS VARGAFTIG\*‡

\*Unité de Pharmacologie Cellulaire, Unité Associée Institut Pasteur–Institut National de la Santé et de la Recherche Médicale 285, Paris, France; and †Unité d’Histopathologie, Institut Pasteur, 25, Rue du Dr. Roux, 75015, Paris, France

Communicated by K. Frank Austen, Harvard Medical School, Boston, MA, September 12, 1995

**ABSTRACT** A murine model for antigen-induced bronchial hyperreactivity (BHR) and airway eosinophilia, two hallmarks of asthma, was developed using ovalbumin-immunized mice, which produce large amounts of IgE (named BP2, “Bons Producteurs 2,” for High Line of Selection 2). A single intranasal ovalbumin challenge failed to modify the bronchial responses, despite the intense eosinophil recruitment into the bronchoalveolar lavage fluid and airways. When mice were challenged twice a day for 2 days or once a day for 10 days, BHR in response to i.v. 5-hydroxytryptamine or to inhaled methacholine was induced in BP2 mice but not in BALB/c mice. Histological examination showed that eosinophils reached the respiratory epithelium after multiple ovalbumin challenges in BP2 mice but remained in the bronchial submucosa in BALB/c mice. Total IgE titers in serum were augmented significantly with immunization in both strains, but much more so in BP2 mice. Interleukin 5 (IL-5) titers in serum and bronchoalveolar lavage fluid of BP2 mice were augmented by the antigenic provocation, and a specific anti-IL-5 neutralizing antibody suppressed altogether airway eosinophilia and BHR, indicating a participation of IL-5 in its development. Our results indicate that the recruitment of eosinophils to the airways alone does not induce BHR in mice and that the selective effect on BP2 mice is related to their increased IgE titers associated with antigen-driven eosinophil migration to the epithelium, following formation and secretion of IL-5.

Bronchial hyperreactivity (BHR), a major characteristic of asthma, is associated with eosinophilic airway inflammation (1–3). By releasing their secretory products, eosinophils are claimed to damage bronchial epithelium and thus to induce BHR (3). This correlation has been demonstrated for guinea pigs (4), but until recently few studies have focused on the development of an eosinophil-dependent BHR model in mice. This may be very important, since such a model would allow exploration of the immunological mechanisms involved in BHR, particularly by taking advantage of the availability of mice strains and of specific tools. We demonstrated that Swiss, CBA, and hypereosinophilic interleukin 5 (IL-5) transgenic mice do not show BHR upon antigenic challenge, despite the marked eosinophil infiltration in airways (5), suggesting the need for factors other than eosinophil infiltration *per se*. The purpose of our study was to develop a model of BHR using a mice selection that produces high antibody titers in response to complex antigens [referred to as BP2, “Bons Producteurs 2”; see Biozzi *et al.* (6)] and that indeed showed very high IgE titers. Following repeated antigenic challenges, these animals became hypereosinophilic and hyperresponsive to nonspecific

bronchoconstrictor substances. Our results suggest that BHR depends on an enhanced IL-5-driven production of eosinophils that locate in the respiratory mucosa, provided the microenvironment contains enough IgE antibodies.

## MATERIALS AND METHODS

**Immunization and Provocation Procedures.** Male BP2 (kindly provided by Denise Mouton, Institut Curie–Centre National de la Recherche Scientifique, Paris and by “Centre d’Elevage R. Janvier”, BP5, 53940 Le Genest Saint-Isle, France) and BALB/c mice (Iffa-Credo and “Centre d’Elevage R. Janvier”), 8–10 weeks of age, were immunized s.c. twice at a 7-day interval with 0.4 ml of a solution of 250 µg of ovalbumin (OA) per ml (Immunobiologicals, Lisle, IL) mixed with 4 mg of Al(OH)<sub>3</sub> per ml (Merck). One week after the second injection, mice were challenged intranasally under light ether anesthesia with 10 µg of OA in 50 µl of 0.9% NaCl (saline) as indicated in *Results*. Control mice were challenged with the vehicle.

**Evaluation of Bronchoconstriction.** In a first procedure, mice were prepared as described (5), using the computerized pulmonary analyzer (Mumed PR800 system, UK) adapted to mice. To evaluate the effect of antigenic challenge on airway responsiveness, 5-hydroxytryptamine (5-HT; Sigma) was injected into the cannulated jugular vein at 10, 20, 40, 80, 160, and 320 µg/kg in a volume of 100 µl during 10 sec at 5-min intervals. The results were expressed as PD30, PD60, and PD90 (the amount of 5-HT needed to augment the bronchial resistance by 30%, 60%, and 90%). In a second procedure, unrestrained conscious mice were placed in a whole body plethysmographic chamber (Buxco Electronics, Sharon, CT) to analyze the respiratory waveforms. After a few minutes for stabilization, an aerosol of methacholine ( $3 \times 10^{-2}$  M in the aerosolator; Aldrich) was delivered during 20 and 60 sec, at a 10-min interval. The airway resistance was expressed as  $Penh = [Te \text{ (expiratory time)}/40\% \text{ of } Tr \text{ (relaxation time)} - 1] \times Pef \text{ (peak expiratory flow)}/Pif \text{ (peak inspiratory flow)} \times 0.67$ , according to the recommendations of the manufacturer. To calculate the  $\Delta Penh$  (difference between the basal and maximal value), the average of five maximal values was used.

**Quantification of Serum IgE Levels.** The total IgE titers were measured by ELISA according to Ledermann *et al.* (7).

**Histological Evaluation.** Lungs were prepared and processed for conventional histological studies. For optic microscopy, the microsections were stained with hematoxylin/eosin and for cyanide-resistant peroxidase (8). For electron micros-

Abbreviations: BALF, bronchoalveolar lavage fluid; BP2, “Bons Producteurs 2”; BHR, bronchial hyperreactivity; OA, ovalbumin; 5-HT, 5-hydroxytryptamine; PD30, PD60, and PD90, doses of 5-HT required to increase airway resistance by 30%, 60%, and 90%; IL, interleukin.

‡To whom reprint requests should be addressed.

copy, ultrathin sections were prepared with an LKB microtome, stained with uranyl acetate/lead citrate, and observed with a JEM-100CXII electron microscope.

For immunohistochemistry, 6- $\mu$ m sections were prepared from OTC-inflated lungs and stained for murine CD4<sup>+</sup> and CD8<sup>+</sup> molecules using rat anti-murine CD4 antibody (L3/T4; Tebu, France) and rat anti-murine CD8 antibody (Ly-2; Tebu, France) and revealed with the alkaline phosphatase anti-alkaline phosphatase technique (9).

**Statistical Analysis.** The significance of differences was determined with the unpaired Student's *t* test.

## RESULTS

**Effect of Immunization and Antigen Challenge on the Cell Composition of the BALF.** Total cell numbers (macrophages essentially) in the BALF of saline-challenged mice amounted to  $0.85 \pm 0.06 \times 10^5$  cells per ml for BP2 and  $1.23 \pm 0.1 \times 10^5$  cells per ml for BALB/c, eosinophils being absent. After a single antigen challenge, these numbers increased time-dependently, peaking at 96 hr (BP2,  $3.95 \pm 0.46$  and  $1.86 \pm 0.24 \times 10^5$  cells per ml; BALB/c,  $2.70 \pm 0.33$  and  $1.03 \pm 0.24 \times 10^5$  cells per ml, for total number and eosinophil number, respectively). These single-challenged mice failed to display BHR (see below) and, accordingly, more intense and/or prolonged stimuli were tested. Mice were challenged either twice a day for 2 days (group indicated as "2 $\times$ 2 days") or once a day for 10 days (group indicated as "1 $\times$ 10 days"). When the BALF was evaluated 24 hr after the last antigen challenge, the cell counts in BALF were increased above those following a single challenge. In particular, the 1 $\times$ 10 days group contained approximately  $4.92 \pm 0.7$  (total cells) and  $3.05 \pm 0.54$  (eosinophils)  $\times 10^5$  cells per ml for BP2 and  $6.02 \pm 0.7$  (total cells) and  $2.99 \pm 0.21$  (eosinophils)  $\times 10^5$  cells per ml for BALB/c, respectively. Lymphocytes were absent in controls but were present in both groups of multichallenged mice, neutrophil counts being unaffected.

**Effect of Antigen Challenge on Bronchial Responsiveness to i.v. 5-HT in Anesthetized Mice.** Single-challenged BP2 mice showed no significant augmentation of bronchial responses to 5-HT (Fig. 1A for 24 hr; other time points not shown). However, BHR was induced following repeated challenge with antigen (Fig. 1A). By contrast, BHR was not uncovered in BALB/c mice (Fig. 1B), even though the eosinophil numbers were comparable to those of BP2. BHR was also absent in response to methacholine in BALB/c mice, no matter what the provocation protocol (data not shown). No statistical correlation (by linear regression analysis) was found between eosinophil numbers in BALF and the intensity of bronchoconstriction in all OA-multichallenged BP2 mice, whether from the groups 2 $\times$ 2 days and 1 $\times$ 10 days.

**Effect of Immunization on the IgE Serum Titers of BALB/c and BP2 Mice.** Nonimmunized BP2 mice contained  $3.35 \pm 2.57$   $\mu$ g of total IgE per ml and BALB/c contained  $0.18 \pm 0.05$   $\mu$ g/ml. Immunization induced a significant augmentation of those titers in both strains, the total amount reaching  $24.96 \pm 5.84$   $\mu$ g/ml and  $0.68 \pm 0.11$   $\mu$ g/ml, in BP2 ( $P < 0.05$ ) and BALB/c ( $P < 0.05$ ) mice, respectively.

**Effect of Antigen Challenge on Bronchial Reactivity to Inhaled Methacholine in Nonanesthetized Mice.** Our inability to induce BHR in BALB/c mice might be attributed to failure of 5-HT to reach the appropriate sites in the airways, since it was injected i.v. A group of immunized and nonanesthetized mice challenged using the protocol 2 $\times$ 2 days was thus evaluated. OA-challenged BP2 mice showed a significant augmentation of airway resistance in response to methacholine as compared to control animals, whereas, once again, no BHR was observed in BALB/c mice (Fig. 2).

**Effect of Immunization and Antigen Challenge on the Lung Histology.** No histological abnormalities were observed in




FIG. 1. Airway responsiveness in immunized and antigen-challenged BP2 (A) and BALB/c (B) mice following different protocols for antigen challenge. In the protocol 1 $\times$ 1 day, 10  $\mu$ g of OA in 50  $\mu$ l of 0.9% NaCl was instilled at  $\approx$ 9 a.m. and BHR was evaluated after 24 hr (see text). In the protocol 2 $\times$ 2 days, OA as above was instilled at 9 a.m. and at 5 p.m.—i.e., at an 8-hr time interval, for 2 days. On the 3rd day, BHR was evaluated. When the protocol 1 $\times$ 10 days was used, OA as above was instilled at 9 a.m. for 10 days. On the 11th day, BHR was evaluated. Controls were saline-challenged mice of each group, which were pooled. Only the BP2 mice challenged 4 (■) or 10 (▨) times showed a significant augmentation ( $\dagger$ ,  $P < 0.001$ , or  $*$ ,  $P < 0.005$ , respectively) of the intensity of bronchoconstriction, compared to control mice. Each value represents the mean  $\pm$  SEM of 5–14 mice.

lungs from control mice of both strains (Fig. 3a). In antigen-challenged mice, granulocytes (mainly eosinophils and a few neutrophils), lymphocytes, plasma cells, and some macrophages accumulated in the peribronchial and perivascular tissues. A moderate submucosal eosinophil infiltration was observed in BP2 and BALB/c mice ( $30 \pm 6$  and  $29 \pm 4$  cells per mm, respectively) challenged once and killed after 24 hr, with practically no eosinophils present in the epithelium. More eosinophils were found in the submucosa of 2 $\times$ 2 days mice ( $120 \pm 7$  cells per mm for BP2 and  $46 \pm 10$  cells per mm for BALB/c). Under those conditions, the epithelium of BALB/c mice was still eosinophil-free, but eosinophils clearly infiltrated that of BP2 mice ( $23 \pm 7$  cells per mm) (Fig. 3b). This epithelial and submucosal eosinophil infiltration in BP2 mice was better appreciated in sections stained for eosinophil peroxidase. Finally, plasma cells were more abundant in the submucosa of BP2 mice (Fig. 3d) as compared to BALB/c mice.

Both mice strains showed a multifocal perivascular and peribronchial eosinophilic alveolitis with eosinophils, macrophages, and few giant cells, even after a single antigenic chal-






FIG. 2. Airway responsiveness in immunized and antigen-challenged BP2 and BALB/c mice in response to inhaled methacholine ( $3 \times 10^{-2}$  M) in the absence of anesthesia. Mice were challenged with OA (■) twice daily for 2 days (protocol 2 $\times$ 2 days) and tested 24 hr after the last challenge. Methacholine was inhaled for 20 and 60 sec at a 10-min interval. The results for 20 sec are shown; similar results were obtained for 60 sec. The results are expressed in  $\Delta$ Penh (see text). Each value represents the mean  $\pm$  SEM of 6–10 mice. Significant differences compared with control (□) are indicated: \*,  $P < 0.05$  and nonsignificance is expressed as N.S.

lenge. Under conditions where lungs from immunized saline-challenged mice contained no T lymphocytes in the bronchial wall, CD4<sup>+</sup> and CD8<sup>+</sup> T lymphocytes were identified in two BP2 mice challenged with the protocol 2 $\times$ 2 days (30 CD4 and 19 CD8 T cells per mm of bronchial wall).

**Electron Microscopy.** Ultrastructural examination confirmed that the majority of the granulocytes observed in the bronchial walls were eosinophils. Eosinophil secondary granules in the epithelium and in the submucosa were of two types, one with a central electron-dense core surrounded by a lighter area and a second showing the reverse—i.e., an electron-dense periphery and a central electron-light core. The morphology of the granules was identical in both strains, with no evidence of degranulation or for differences in number of granules. Thus, an average of  $22.7 \pm 1.7$  and  $19.44 \pm 3$  granules per eosinophil section of BP2 mice was scored in the submucosa ( $n = 18$  cells) and in the epithelium ( $n = 9$  cells), respectively. Similar values were obtained for eosinophils of BALB/c mice counted in the submucosa.

**Interference of the Anti-IL-5 Antibody TRFK-5 with Eosinophilia and BHR.** Approximately 150 pg of IL-5 per ml was found in the serum and 300 pg/ml in the BALF of BP2 mice 1–6 hr after the fourth antigenic provocation (protocol 2 $\times$ 2 days). This led us to evaluate the anti-IL-5 antibody TRFK-5. In these experiments, OA challenges augmented the total cell numbers in the BALF of IgG isotype-treated mice to  $6.3 \pm 0.8 \times 10^5$  cells per ml ( $P < 0.001$ ), a number comparable to that of saline-treated animals (Fig. 4A). This increment was accounted for by a marked increase in the proportion of eosinophils ( $57.5\% \pm 2.5\%$  of total cell numbers). TRFK-5, injected i.v. at 0.5 mg per mouse 1 hr before each OA challenge, reduced significantly the total BALF cell counts, which were accounted for essentially by decreased eosinophil numbers (isotype,  $3.54 \pm 0.38 \times 10^5$  cells per ml; TRFK-5,  $0.58 \pm 0.25 \times 10^5$  cells per ml;  $P < 0.05$ ) (Fig. 4A). Histological observations confirmed that the administration of TRFK-5 inhibited the epithelial (from 6.8 to 0.7 cells per mm) and submucosal (from 55.7 to 14.2 cells per mm) eosinophil infiltration in two mice.

As expected, OA challenges in saline or IgG isotype-treated mice resulted in BHR (Fig. 4B). TRFK-5 inhibited the antigen-induced BHR for 5-HT in anesthetized (Fig. 4B) and for methacholine in nonanesthetized antigen-challenged mice (Fig. 4C).

## DISCUSSION

Eosinophils are recruited to the airways of asthmatic patients and are considered important effector cells in BHR, a hall-




FIG. 4. Effect of the anti-IL-5 antibody TRFK-5 on eosinophil recruitment into the BALF (A) and on BHR (B and C) in immunized BP2 mice. The protocol 2 $\times$ 2 days was used. Purified IL-5 antibody from TRFK-5 cells was injected i.v. at 0.5 mg per mouse to immunized BP2 mice, 1 hr before each of the four OA (ovo) challenges. The same volume of control antibody (rat IgG) was injected into control mice. Bronchoalveolar lavage and BHR were evaluated 24 hr after the last antigen challenge. BHR with 5-HT was expressed as PD90 (see text) in anesthetized mice and BHR with methacholine was expressed as  $\Delta$ Penh (see text) in nonanesthetized mice. Each value represents the mean  $\pm$  SEM of 4–12 mice. Significant differences compared with control isotype (challenged with OA) are indicated: \*,  $P < 0.05$ ; †,  $P < 0.005$ ; ‡,  $P < 0.001$ .

mark of asthma (10). Although BHR is reported to correlate with the number of eosinophils in lungs staining positively with a monoclonal antibody that recognizes secreted eosinophil cationic protein (10–13), there is no formal evidence for a causal relationship between eosinophil recruitment *per se* and BHR in humans (14, 15), sheep (16), and guinea pigs (4). Few studies have addressed the correlation between eosinophil recruitment to the airways and BHR in mice, possibly because the evaluation of their pulmonary functions was considered difficult or unreliable. Nevertheless, murine models should be invaluable for dissecting the mechanisms of BHR, since murine basic immunology is well defined. This led us to search for and to identify mice that displayed intense anaphylactic responses, in addition to antigen-induced hyper eosinophilia. We took advantage of the availability of high-responder mice (detailed in ref. 6) selected by bidirectional breeding for antibody responsiveness to sheep erythrocytes in terms of agglutinin titers, which have been shown to be homozygous for the character considered in F<sub>14</sub>–F<sub>17</sub> generations (6, 17). These mice developed BHR, which was detected by two methods. In




FIG. 3. (a) Lobar bronchus from BP2 mice immunized with OA and challenged with saline. Control BALB/c mice showed the same morphology as BP2 mice. Note the absence of inflammatory cell infiltration. ep, Epithelium; mus, bronchial muscle; sm, submucosa; alv, alveolus. ( $\times 70$ .) (b) Lobar bronchus from BP2 mice challenged with OA according to protocol 2 $\times$ 2 days. A marked eosinophil infiltration of the submucosa (three short arrows) and of the mucosa (three arrowheads) is noted. ( $\times 70$ .) (c) Higher magnification of the section in b showing the mucosa containing numerous infiltrating eosinophils (two arrowheads). ( $\times 280$ .) (d) Higher magnification of the mucosa and submucosa showing the dominant infiltrating eosinophils (short arrows) but also a number of plasma cells (long arrows). ( $\times 280$ .) All sections were stained with hematoxylin/eosin.

a first one, 5-HT was administered i.v. to anesthetized mice (5, 18), and in the second, methacholine was administered by aerosol to awake animals (19). This is important since bronchoconstrictor responses may differ according to the route of administration of the agonists, which determines the first line cell species that is encountered. Furthermore, since eosinophils activated *in situ* might induce BHR by affecting the respiratory epithelium (3), it seemed logical to deliver the bronchoconstrictor agent directly into the airways. Both methods provided similar results.

BP2 mice differed from BALB/c mice in at least four different ways. (i) A first and essential difference is that only BP2 mice displayed BHR following multiple antigenic challenges. (ii) They showed very high levels of serum IgE, which doubled after immunization, whereas BALB/c mice had considerably lower amounts, which, as expected, also augmented upon immunization. (iii) BP2 mice underwent an intense anaphylactic bronchoconstriction when challenged with i.v. OA, whereas BALB/c were unresponsive (not shown). (iv)

Eosinophils were identified in the epithelium of BP2 mice after antigen challenges, being absent in BALB/c mice. In previous studies, we also failed to identify eosinophils in the epithelium of antigen-challenged mice from other strains, including IL-5 transgenic animals, even though the BALF and the submucosa were heavily enriched with eosinophils (5). The basic difference between BP2 and other mice in terms of eosinophils thus appears to relate to their topographic distribution and not to absolute counts. It was tempting to speculate that eosinophils stimulated at the vicinity of the targeted respiratory epithelium, in a hyper-IgE environment, are likely to become activated, particularly after repeated antigenic challenges. Nevertheless, the secondary granules of the eosinophils found in the epithelium and in the submucosa did not show morphological evidence of degranulation, according to published criteria (20, 21). It is thus unlikely that eosinophils participate in the induction of BHR in BP2 mice, via the secretion of cationic proteins such as "Major Basic Protein" as in guinea pigs (4), and it also remains to be shown whether an augmented

production of lipid mediators may occur. Eosinophils from immunized and aerosol-challenged guinea pigs are indeed primed to produce larger amounts of leukotriene C<sub>4</sub> than control animals (22). On the other hand, the respiratory epithelium might express adhesion proteins in the hyper-IgE environment in BP2 mice, leading to the recruitment and activation of eosinophils. Indeed, we demonstrated that an antibody against very late activation antigen 4 suppresses the antigen-induced airway eosinophil recruitment and *in situ* activation and the associated BHR in guinea pigs (23).

Our results contrast to those of Renz *et al.* (18), who showed that BALB/c mice immunized by inhaled OA for a long time interval display BHR in response to i.v. methacholine. This may result from different protocols, particularly since no eosinophils or other inflammatory cells were found in the airways of these mice at the time of BHR but might have occurred before. BHR persists in BP2 mice for at least 16 days, a time when eosinophils are absent (not shown).

In immunized BP2 mice, the IgE titers were very elevated and the number of plasma cells identified in the bronchial submucosa was increased, two possibly related events. Increased IgE titers may be involved in the induction of BHR, particularly since an anti-IgE antibody inhibits altogether airway eosinophilia and BHR in our model (data not shown). IgE production is dependent on IL-4 (24), a Th2 lymphocyte-derived cytokine. IL-4 is also involved in eosinophil recruitment to the airways upon antigen challenge (25), but it is clear that IL-5 is responsible for eosinophil recruitment to the airways in the mice strains studied and for BHR, when applicable. Indeed, IL-5 was present in the BALF and serum of the BP2 mice after antigen challenge and the anti-IL-5 antibody TRFK-5 suppressed altogether eosinophilia in airways and tissues and BHR. In allergen-challenged guinea pigs, anti-IL-5 antibody also suppressed airway eosinophilia (26) and BHR (27, 28). CD4<sup>+</sup> T lymphocytes that migrate into the lungs after antigen challenge (29) are responsible for the local production of IL-4 and IL-5, which are important in B- and T-lymphocyte development and function and in eosinophil activation and differentiation (30). In CD4<sup>+</sup> deficient or immune-depleted mice, BALF eosinophilia and airway inflammation in response to antigen provocation were absent (25, 31) and BHR was reduced (25). In confirmation, we identified CD4<sup>+</sup> T cells in the bronchial wall after antigen challenge in BP2 mice, suggesting that these cells participate in the induction of airway eosinophilia, and accordingly of BHR, by releasing cytokines.

As a conclusion, repeated intranasal antigenic challenges induce eosinophil recruitment to the lungs and to the BALF in BALB/c and in BP2 mice, but BHR develops selectively in the latter. Since eosinophil recruitment to the lungs and BHR were inhibited by *in vivo* IL-5 neutralization, eosinophils are clearly necessary, but not sufficient, for BHR. Our finding that eosinophils locate in the respiratory epithelium only in hyper-IgE BP2 mice, which become hyperresponsive, suggests that BHR involves the interaction between eosinophils recruited and/or primed by IL-5 and a component of the epithelium in the presence of an altered microenvironment, including high IgE titers. Interference with one of the sequential events involved—i.e., excessive eosinophil production or recruitment, the selective expression of adhesion proteins on the endothelial or epithelial surfaces, the production or the effects of IgE, or finally the production or effects of eosinophil-borne mediators—should be sufficient to suppress BHR.

We are grateful to Drs. Christopher Heusser and Jürgen Koepper (CIBA-Geigy, Basel, Switzerland) for the IgE determinations, to Drs.

Marina Pretolani and Michel Chignard for the critical review of the manuscript, and to Mr. R. Janvier (Centre d'Élevage R. Janvier, France) for help and support in producing BP2 mice. Dr. Antonio Vinas de Souza participated in the early part of our work.

1. Boushey, H. A., Holtzman, M. J., Sheller, J. R. & Nadel, J. A. (1980) *Am. Rev. Respir. Dis.* **121**, 389–413.
2. Corrigan, C. J. & Kay, A. B. (1992) *Immunol. Today* **13**, 501–507.
3. Gleich, G. J. (1990) *J. Allergy Clin. Immunol.* **85**, 422–436.
4. Pretolani, M., Ruffié, C., Joseph, D., Campos, M. G., Church, M. K., Lefort, J. & Vargaftig, B. B. (1994) *Am. J. Respir. Crit. Care Med.* **149**, 1167–1174.
5. Lefort, J., Bachelet, C. M., Leduc, D. & Vargaftig, B. B. (1995) *J. Allergy Clin. Immunol.*, in press.
6. Biozzi, G., Mouton, D., Sant'Anna, O. A., Passos, H. C., Gennari, M., Reis, M. H., Ferreira, V. C. A., Heumann, A. M., Bouthillier, Y., Ibanez, O. M., Stiffel, C. & Siqueira, M. (1979) *Curr. Top. Microbiol. Immunol.* **85**, 31–98.
7. Ledermann, F., Schlienger, C., Wagner, K. & Heusser, Ch. (1991) *J. Immunol. Methods* **141**, 263–275.
8. Ten, R. M., Pease, L. R., McKean, D. J., Bell, M. P. & Gleich, G. J. (1989) *J. Exp. Med.* **169**, 1757–1769.
9. Polak, J. M. & Noorden, S. V. (1986) in *Immunocytochemistry: Modern Methods and Applications* (Wright, Bristol, U.K.), 2nd Ed., pp. 47–48.
10. Djukanovic, R., Roche, W. K., Wilson, J. W., Beasley, C. R. W., Twentyman, O. P., Howarth, P. H. & Holgate, S. T. (1990) *Am. Rev. Respir. Dis.* **142**, 434–457.
11. Howarth, P. H., Bradding, P., Montefort, S., Peroni, D., Djukanovic, R., Carroll, M. P. & Holgate, S. T. (1994) *Am. J. Respir. Crit. Care Med.* **150**, S18–S22.
12. Azzawi, M., Bradely, B., Jeffery, P. K., Frew, A. J., Wardlaw, A. J., Knowles, G., Assoufi, B., Collins, J. V., Durham, S. & Kay, A. B. (1990) *Am. Rev. Respir. Dis.* **142**, 1407–1413.
13. Tai, P. C., Spry, C. J. F., Peterson, C., Venge, P. & Olsson, I. (1984) *Nature (London)* **309**, 182–184.
14. Gleich, G. J., Frigas, E., Loegering, D. A., Wassom, D. L. & Steinmuller, D. (1979) *J. Immunol.* **123**, 2925–2927.
15. Frigas, E., Loegering, D. A. & Gleich, B. S. (1980) *Lab. Invest.* **42**, 35–43.
16. Abraham, W. M., Sielczak, M. W., Ahmed, A., Cortes, A., Lauredo, I. T., Kim, J., Pepinsky, B., Benjamin, C. D., Leone, D. R., Lobb, R. R. & Weller, P. F. (1994) *J. Clin. Invest.* **93**, 776–787.
17. Mouton, D., Siqueira, M., Sant'Anna, O. A., Bouthillier, Y., Ibanez, O., Ferreira, V. C. A., Mevel, J. C., Reis, M. H., Piatti, R. M., Stiffel, C. & Biozzi, G. (1988) *Eur. J. Immunol.* **18**, 41–49.
18. Renz, H., Smith, H. R., Henson, J. E., Ray, S., Irvin, C. G. & Gelfand, E. W. (1992) *J. Allergy Clin. Immunol.* **89**, 1127–1138.
19. Chand, N., Nolan, K., Pillar, J., Lomask, M., Diamantis, W. & Sofia, R. D. (1992) *Allergy* **48**, 230–235.
20. Beasley, R., Roche, W. R., Roberts, J. A. & Holgate, S. T. (1989) *Am. Rev. Respir. Dis.* **139**, 806–817.
21. Dvorak, A. M. (1994) in *Eosinophils in Allergy and Inflammation*, eds. Gleich, G. J. & Kay, A. B. (Marcel, New York), pp. 159–209.
22. Pretolani, M., Joseph, D., Lobb, R. R. & Vargaftig, B. B. (1994) *Am. J. Respir. Crit. Care Med.* **151**, A221.
23. Pretolani, M., Ruffié, M., Lapa e Silva, J. R., Joseph, D., Lobb, R. R. & Vargaftig, B. B. (1994) *J. Exp. Med.* **180**, 795–805.
24. Kopf, M., Le Gros, G., Bachmann, M., Lamers, M. C., Bluethmann, H. & Köhler, G. (1993) *Nature (London)* **362**, 245–248.
25. Brusselle, G. G., Kips, J. C., Tavernier, J. H., Van Der Heyden, J. G., Cuvelier, C. A., Pauwels, R. A. & Bluethmann, H. (1994) *Clin. Exp. Allergy* **24**, 73–80.
26. Chand, N., Nolan, K., Pillar, J., Lomask, M., Diamantis, W. & Sofia, R. D. (1992) *Allergy* **48**, 230–235.
27. Nakajima, H., Iwamoto, I., Tomoe, S., Matsumura, R., Tomioka, H., Takatsu, K. & Yoshida, S. (1992) *Am. Rev. Respir. Dis.* **146**, 374–377.
28. Van Oosterhout, A. J. M., Ladenius, A. R. C., Savelkoul, H. F. J., Ark, I. V., Delsman, K. C. & Nijkamp, F. P. (1993) *Am. Rev. Respir. Dis.* **147**, 548–552.
29. Gerblich, A. A., Salik, H. & Schuyler, M. R. (1991) *Am. Rev. Respir. Dis.* **143**, 533–537.
30. Garlich, C. G., Falcone, A., Kung, T. T., Stelts, D., Penline, K. J., Beavis, A. J., Smith, S. R., Egan, R. W. & Umland, S. P. (1995) *Clin. Immunol. Immunopathol.* **75**, 75–83.
31. Gavett, S. H., Chen, X., Finkelman, F. & Wills-Karp, M. (1994) *Am. J. Respir. Crit. Care Med.* **10**, 587–593.

Attorney Docket No.: 5051-451ip

PATENT

**IN THE UNITED STATES PATENT AND TRADEMARK OFFICE**

In Re: Yuehua Li, et al.

Confirmation No.: 8515

Serial No.: 09/914,020

Group Art Unit: 1635

Filed: December 31, 2001

Examiner: Janet L. Epps-Ford

For: METHODS AND COMPOSITIONS FOR ALTERING MUCOUS SECRETION

**Exhibit 3**

**EXHIBIT 3****Mucin Secretion in a Murine Model of Asthma**

<b>Peptide</b>	<b>% Mucin Secretion</b>	<b>% Inhibition of Mucin Secretion Relative to OVA Control</b>
<b>Untreated Saline Control</b>	<b>8</b>	<b>92</b>
<b>OVA Control</b>	<b>100</b>	<b>0</b>
<b>BIO-124R (RNS)</b>	<b>116</b>	<b>-16</b>
<b>BIO-124 (MANS)</b>	<b>24</b>	<b>76</b>
<b>BIO-120</b>	<b>23</b>	<b>77</b>
<b>BIO-116</b>	<b>32</b>	<b>68</b>
<b>BIO-112</b>	<b>23</b>	<b>77</b>
<b>BIO-110</b>	<b>34</b>	<b>66</b>